



## King's Research Portal

DOI:

[10.1186/s13550-017-0256-5](https://doi.org/10.1186/s13550-017-0256-5)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Nair, A., Veronese, M., Xu, X., Curtis, C., Turkheimer, F., Howard, R., & Reeves, S. (2017). Erratum to Test-retest analysis of a non-invasive method of quantifying [(11)C]-PBR28 binding in Alzheimer's disease. *EJNMMI Research*, 7(1), 14. <https://doi.org/10.1186/s13550-017-0256-5>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

ERRATUM

Open Access



# Erratum to: Test-retest analysis of a non-invasive method of quantifying [(11)C]-PBR28 binding in Alzheimer's disease

Akshay Nair<sup>1\*</sup>, Mattia Veronese<sup>2</sup>, Xiaohui Xu<sup>3,4</sup>, Charles Curtis<sup>3,4</sup>, Federico Turkheimer<sup>2</sup>, Robert Howard<sup>5</sup> and Suzanne Reeves<sup>5</sup>

## Erratum

Following publication of the original article [1] the authors contacted the team with the request that some additional information be added to the article. Please see below for details:

### Additional Table: absolute variability

Variability (VAR) presented in the main article represents **mean variability** between test and retest, not absolute variability as stated. The authors would like to apologise for this error. The absolute variability is presented in the table below has been calculated in the following manner:

$$\text{Absolute variability} = \frac{2 * (|\text{Test} - \text{Retest}|)}{(\text{Test} + \text{Retest})} * 100$$

These analyses allow for easier comparison with other studies however do not alter the findings or discussion of the paper – SUV normalised to whole brain (SUV<sub>WB</sub>) shows the lowest variability in our cohort, with increased variability found in the unadjusted SUV and the SUV normalised to cerebellum. As compared to other test-retest studies discussed in the paper, the variability for SUV based methods remains favourable in this pilot cohort. These results should be validated in larger studies.

## Author details

<sup>1</sup>UCL Huntington's Disease Research Group, 2nd floor, Russell Square House, 10-12 Russell Square, London, WC1B 5EH, UK. <sup>2</sup>Centre for Neuroimaging Sciences, IoPPN, King's College London, Box PO89De Crespigny Park, London SE5 8AF, UK. <sup>3</sup>MRC Social, Genetic and Developmental Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, De Crespigny Park, Denmark Hill, London SE5 8EF, UK. <sup>4</sup>NIHR Biomedical Research Centre for Mental Health, Maudsley Hospital and Institute of Psychiatry, Psychology & Neuroscience, King's College London, London SE5 8AF, UK. <sup>5</sup>Division of Psychiatry, University College London, 6th Floor, Wings A and B, Maple House, 149 Tottenham Court Road, London W1T 7NF, UK.

\* Correspondence: akshay.nair@ucl.ac.uk

<sup>1</sup>UCL Huntington's Disease Research Group, 2nd floor, Russell Square House, 10-12 Russell Square, London, WC1B 5EH, UK

Received: 5 January 2017 Accepted: 5 January 2017

Published online: 14 February 2017

## References

1. Nair A, et al. Test-retest analysis of a non-invasive method of quantifying [(11)C]-PBR28 binding in Alzheimer's disease. *EJNMMI Res.* 2016;6:72.

	Raw SUV	SUV <sub>WB</sub>	SUV <sub>C</sub>
Frontal Lobe	6.05 (2.45)	1.01 (0.67)	4.36 (4.84)
Parietal Lobe	6.69 (2.33)	0.97 (0.77)	5.15 (5.10)
Temporal Lobe	5.92 (3.13)	0.61 (0.49)	3.78 (4.92)
Occipital Lobe	6.81 (2.83)	1.07 (0.88)	3.87 (3.79)
Hippocampus	5.82 (1.95)	1.57 (1.34)	5.35 (6.50)
Parahippocam-pal cortex	5.05 (3.78)	2.07 (2.58)	5.77 (7.96)
Posterior Cingulate	6.58 (2.60)	1.89 (1.33)	5.92 (5.97)
Amygdala	3.45 (4.50)	4.31 (3.17)	6.54 (8.56)
Cerebellum	7.18 (7.84)	4.18 (5.07)	
Whole Brain	6.09 (2.73)		
Mean (SD)	5.96 (3.41)	1.96 (1.81)	5.09 (5.95)